






















Infants', Children's and Jr. **TYLENOL**[®]

Dosing Information for Healthcare Professionals

Use this chart to determine the proper dose of **TYLENOL**[®] for your patient. **If possible, use weight to dose; otherwise use age.**

DOSE — Give your patient the dose indicated below every 4 hours as needed. Do not give more than 5 doses in 24 hours.

mL = milliliter
tsp = teaspoonful

Weight (lbs)	Age	 Infants' TYLENOL[®] Oral Suspension Active Ingredient: Acetaminophen 160mg (in each 5mL or 1 tsp) Use only as directed.	 Children's TYLENOL[®] Oral Suspension Active Ingredient: Acetaminophen 160mg (in each 5mL or 1 tsp) Use only as directed.	 Children's TYLENOL[®] Meltaway Chewable Tablets Active Ingredient: Acetaminophen 80mg (in each tablet) Use only as directed.	 Jr. TYLENOL[®] Meltaway Chewable Tablets Active Ingredient: Acetaminophen 160mg (in each tablet) Use only as directed.
6-11 lbs	0-3 mos	1.25 mL 	—	—	—
12-17 lbs	4-11 mos	2.5 mL 	—	—	—
18-23 lbs	12-23 mos	3.75 mL 	—	—	—
24-35 lbs	2-3 yrs	5 mL 	5 mL (1 tsp) 	2 tablets 	—
36-47 lbs	4-5 yrs	—	7.5 mL (1½ tsp) 	3 tablets 	—
48-59 lbs	6-8 yrs	—	10 mL (2 tsp) 	4 tablets 	2 tablets 
60-71 lbs	9-10 yrs	—	12.5 mL (2½ tsp) 	5 tablets 	2½ tablets 
72-95 lbs	11 yrs	—	15 mL (3 tsp) 	6 tablets 	3 tablets 

Remind parents and caregivers to:

- Read and follow the label on all **TYLENOL**[®] products
- Take every 4 hours as needed
- Do NOT exceed more than 5 doses in 24 hours
- Do NOT use with any other product containing acetaminophen
- Keep all medicines out of the reach of children
- Do NOT administer adult medicines to children
- Use only the dosing device that comes with a specific product:
- Infants' **TYLENOL**[®] Oral Suspension — enclosed SimpleMeasure[™] syringe
- Children's **TYLENOL**[®] Oral Suspension — enclosed measuring cup

- Children's **TYLENOL**[®] Meltaway Chewable Tablets are not the same concentration as Jr. Strength **TYLENOL**[®] Meltaway Chewable Tablets. Jr. **TYLENOL**[®] Meltaway Chewable Tablets contain 160mg of acetaminophen, while Children's **TYLENOL**[®] Meltaway Chewable Tablets contain 80mg of acetaminophen
- All Infants' and Children's **TYLENOL**[®] Oral Suspension products in stores have the same acetaminophen concentration (160mg/5mL)

TYLENOL[®]
ACETAMINOPHEN



Children's Advil[®] Relief you can trust

Children's Advil[®]



Measure dosing only with the device that comes with the product.



Available in 4 flavors!
 • Dye-Free
 • White Grape
 • Fruit
 • Grape
 • Blue Raspberry



Infants' Advil[®] Concentrated Drops
for ages 6-23 months

Children's Advil[®] Suspension
for ages 2-11 years

Junior Strength Advil[®] Chewables
for ages 6-11 years

Junior Strength Advil[®] Tablets
for ages 6-11 years

Advil[®] Tablets Advil[®] Liqui-Gels[®]
for ages 12 and up

Active ingredient

Ibuprofen 50 mg (NSAID)¹
in each 1.25 mL

Ibuprofen 100 mg (NSAID)¹
in each 5 mL

Ibuprofen 100 mg (NSAID)¹
in each tablet

Ibuprofen 100 mg (NSAID)¹
in each tablet

Ibuprofen 200 mg (NSAID)¹
in each tablet/capsule

Indications

Fever reducer, relieves minor aches and pains due to the common cold, flu, headaches, and toothaches

Fever reducer, relieves minor aches and pains due to the common cold, flu, sore throat, headaches, and toothaches

Fever reducer, relieves minor aches and pains due to the common cold, flu, sore throat, headaches, and toothaches

Fever reducer, relieves minor aches and pains due to the common cold, flu, sore throat, headaches, and toothaches

Fever reducer, relieves minor aches and pains due to headache, toothache, backache, menstrual cramps, the common cold, muscular aches, and minor pain of arthritis

Weight Age

under 6 months

Do not use

Do not use

Do not use

Do not use

Do not use

12-17 pounds 6-11 months

1.25 mL¹ every 6-8 hrs not to exceed 4 doses/24 hrs

Use Infants' Advil[®] Concentrated Drops

Use Infants' Advil[®] Concentrated Drops

Use Infants' Advil[®] Concentrated Drops

Use Infants' Advil[®] Concentrated Drops

18-23 pounds 12-23 months

1.875 mL¹ every 6-8 hrs not to exceed 4 doses/24 hrs

↓

↓

↓

↓

24-35 pounds 2-3 years

Use Children's Advil[®] Suspension

1 tsp¹ every 6-8 hrs not to exceed 4 doses/24 hrs

Use Children's Advil[®] Suspension

Use Children's Advil[®] Suspension

Use Children's Advil[®] Suspension

36-47 pounds 4-5 years

↓

1½ tsp¹ every 6-8 hrs not to exceed 4 doses/24 hrs

↓

↓

↓

48-59 pounds 6-8 years

Use Children's Advil[®] Suspension or Junior Strength Advil[®] Chewables or Tablets

2 tsp¹ every 6-8 hrs not to exceed 4 doses/24 hrs

2 tablets every 6-8 hrs not to exceed 4 doses/24 hrs

2 tablets every 6-8 hrs not to exceed 4 doses/24 hrs

↓

60-71 pounds 9-10 years

↓

2½ tsp¹ every 6-8 hrs not to exceed 4 doses/24 hrs

2½ tablets every 6-8 hrs not to exceed 4 doses/24 hrs

2 tablets every 6-8 hrs not to exceed 4 doses/24 hrs

↓

72-95 pounds 11 years

↓

3 tsp¹ every 6-8 hrs not to exceed 4 doses/24 hrs

3 tablets every 6-8 hrs not to exceed 4 doses/24 hrs

3 tablets every 6-8 hrs not to exceed 4 doses/24 hrs

↓

12+ years

Use Advil[®] Tablets or Liqui-Gels[®]

Use Advil[®] Tablets or Liqui-Gels[®]

Use Advil[®] Tablets or Liqui-Gels[®]

Use Advil[®] Tablets or Liqui-Gels[®]

1 or 2 tablets/capsules¹ every 4-6 hrs not to exceed 6 tablets/capsules/24 hrs

¹ 1 tsp=100 mg ibuprofen per 5 mL

² Nonsteroidal anti-inflammatory drug.

³ Measure dosing only with the device provided.

⁴ If pain or fever does not respond to 1 tablet/capsule, 2 tablets/capsules may be used.

2 tablets/capsules may be used.

Shake liquid products well before dosing.

Please refer to Drug Facts on the product label or visit advil.com for complete warnings.

Important Instructions for Proper Use

- Read and keep the cartons for complete warnings and information on Advil[®] products
- Remind patients that if they are taking other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), they should consult with you before taking any Advil[®] products
- Adult strength medicines should not be administered to children