

IBUPROFEN Dosing Chart

DO NOT USE FOR CHILDREN UNDER 6 MONTHS OF AGE UNLESS DIRECTED BY PHYSICIAN.

ml= milliliter mg= milligrams tsp = teaspoon	Infant Drops 50 mg/ 1.25 mL	Liquid 100 mg/ 5 mL (tsp)	Chewable Tablets 50 mg in each tablet	Junior Strength Chewable Tablets 100 mg in each tablet	Adult 200 mg in each tablet
Weight (in pounds)	Dose	Dose	Dose	Dose	Dose
12-17 pounds	1.25 ml	½ tsp	—	—	—
18-23 pounds	1.875 ml	¾ tsp	—	—	—
24-35 pounds	2.50 ml	1 tsp	2 tablets	—	—
36-47 pounds	3.75 ml	1½ tsp	3 tablets	—	—
48-59 pounds	5.00 ml	2 tsp	4 tablets	2 tablets	1 tablet
60-71 pounds	—	2½ tsp	5 tablets	2½ tablets	1 tablet
72-95 pounds	—	3 tsp	6 tablets	3 tablets	1½ tablets
96+ pounds	—	4 tsp	8 tablets	4 tablets	2 tablets

Remember

- repeat dosage every 6-8 hours as needed or directed by physician
- do not use with any other product containing ibuprofen; do not use multi-ingredient products in children under 6 years of age
- check with your healthcare provider if you have any questions
- check with your healthcare provider when giving medication to infants under 12 pounds or children under 2 years of age
- always read and follow the label directions
- whenever possible, use weight to determine dosage
- for accurate dosing, only use the dosing device that comes with the product
- do not give adult medicines to infants/children
- keep all medicines out of the reach of children