

Overcoming Anchorage Limitations

TADs

Temporary anchorage devices (TADs) for orthodontic anchorage are widely accepted. They are changing the way orthodontists treat some patients' malocclusions. TADs provide a fixed point from which to apply force to move teeth. They can be placed in many different sites in the mouth. Placement is customized for each patient. TADs may contribute to predictable results, shorter treatment time and completion of active treatment on schedule.



TADs range from 6 to 12 millimeters in length.

There is little or no discomfort when a TAD is placed. Caring for your TAD generally requires only routine brushing.

Although small, the TAD does a big job of helping your orthodontist move your teeth predictably into their optimal positions.

Your orthodontist, a member of the American Association of Orthodontists, sincerely thanks you for placing your confidence in him/her to treat your orthodontic needs.

Orthodontists receive an additional two to three years of specialized education beyond dental school to learn the proper way to align and straighten teeth. Only those with this education may call themselves "orthodontists," and only orthodontists are eligible for membership in the American Association of Orthodontists.

Photos/illustrations are used with permission by Under Dog Media, LP from OrthoTADs: The Clinical Guide and Atlas, www.orthotads.com

American Association of Orthodontists
401 North Lindbergh Boulevard
St. Louis, MO 63141-7816

314.993.1700
800.STRAIGHT (consumer information line)

www.braces.org

Email: info@aaortho.org

American Association of
Orthodontists



© 2008 American Association of Orthodontists

Temporary Anchorage Devices (TADs)

For Predictable Tooth Movement




Questions and Answers

What are TADs (temporary anchorage devices)?

TADs are titanium-alloy mini-screws, ranging from 6 to 12 millimeters in length and 1.2 to 2 millimeters in diameter. They are fixed to bone temporarily to enhance orthodontic anchorage. Titanium alloys have been used as joint replacements and for dental implants for many years. These alloys are not rejected by the body.

Orthodontists often use holding arches, also known as space maintainers, and headgear to control anchorage and minimize the movement of certain teeth while carrying out the desired movement of other teeth. TADs allow orthodontists to overcome limitations of holding arches and headgear and perform difficult tooth movements predictably.



A TAD is inserted.

TADs can also provide a point of anchorage for patients with missing teeth.

Where are TADs placed?

TADs are placed in the bone between the roots of the teeth and can be placed in the bone in the roof of the mouth as well.

A Small Device That Does a Big Job

How is a TAD placed and does it hurt?

Placement is minimally invasive and often completed using only topical anesthetic. TADs are inserted directly into the bone using a special instrument. There is little or no discomfort because there are no nerve endings in the bone tissue. Once placed, the orthodontist is able to use the TAD as orthodontic anchorage immediately.

Because of the possibility that TADs can loosen or fall out, patients should avoid picking or pulling at the TAD. If the TAD does become loose or come out, call your orthodontist as soon as possible.



The site for the TAD is selected.



The TAD immediately after placement.

Who places the TAD?

Because orthodontists have the training and expertise to place them, many orthodontists place TADs themselves. This ensures the TAD is placed exactly where the orthodontist wants it. Some orthodontists may choose to have a TAD placed by another dental specialist.



A TAD serves as an anchor to perform difficult tooth movements predictably.

How long are TADs left in?

Your orthodontist will advise you about how long a TAD will be needed. It may be required only for a few months, or it may be needed throughout your orthodontic treatment. TADs are versatile and may be used in different areas of the mouth during different parts of treatment.

Can TADs be used for patients at any age?

TADs are placed on patients who have permanent teeth. Patients with active periodontal (gum) disease may not be candidates for TADs.