

## COMMON MEDICATION DOSING

### Acetaminophen (Tylenol) Dosage Chart

Give every 4 to 6 hours if needed for fever or pain

Weight (Pounds)	Dose (mgs)	Infant Oral Suspension (160mg/5ml)	Children's Oral Suspension (160mg.ml)	Chewable (160mg)	Adult Tablet (325mg)
6-11 lbs	40 mg	1.25 ml	---	---	---
12-17 lbs	80 mg	2.5 ml	2.5 ml	---	---
18-23 lbs	120 mg	3.75 ml	3.75 ml	---	---
24-35 lbs	160 mg	---	5 ml	1 tablet	---
36-47 lbs	240 mg	---	7.5 ml	1 ½ tablets	---
48-59 lbs	320 mg	---	10 ml	2 tablets	1 tablet
60-71 lbs	400 mg	---	12.5 ml	2 ½ tablets	1 tablet
72-95 lbs	480 mg	---	15 ml	3 tablets	1 ½ tablets
96+ lbs	640 mg	---	20 ml	4 tablets	2 tablets

### Ibuprofen Dosage Chart

Give every 6 hours if needed for fever or pain. Do NOT give to children under 6 months of age

Weight (Pounds)	Dose (mgs)	Infant Drops (50mg/1.25ml)	Children's Liquid (100mg/5ml)	Chewable Tablets (50mg)	Junior Strength Tablets (100mg)	Adult Tablet (200mg)
6-11 lbs	---	---	---	---	---	---
12-17 lbs	50 mg	1.25 ml	2.5 ml	---	---	---
18-23 lbs	75 mg	1.875 ml	3.75 ml	1 ½ tablets	---	---
24-35 lbs	100 mg	2.5 ml	5 ml	2 tablets	1 tablet	---
36-47 lbs	150 mg	---	7.5 ml	3 tablets	1 ½ tablets	---
48-59 lbs	200 mg	---	10 ml	4 tablets	2 tablets	1 tablet
60-71 lbs	250 mg	---	12.5 ml	5 tablets	2 ½ tablets	1 tablet
72-95 lbs	300 mg	---	15 ml	6 tablets	3 tablets	1 ½ tablets
96+ lbs	400 mg	---	20 ml	8 tablets	4 tablets	2 tablets

### Diphenhydramine (Benadryl) Dosage Chart

Give every 6 hours if needed

Weight (Pounds)	Liquid (12.5mg/5ml)	Chewable (12.5mg)	Tablets or Capsules (25mg)
20-24 lbs	4 ml	---	---
25-37 lbs	5 ml	1 tablet	½ tablet
38-49 lbs	7.5 ml	1 ½ tablets	½ tablet
50-99 lbs	10 ml	2 tablets	1 tablet
100+ lbs	---	4 tablets	2 tablets