

## 351014 ROSACEA CREAM

### What is this preparation?

---

This preparation is custom-formulated specifically for you to better treat your individual needs. The unique combination of active and inactive ingredients has been selected to produce an outcome that is clinically different to that which is commercially available. For this reason, no standardized information or literature is available for your final preparation. Custom-created preparations allow us to meet your individual needs without being restricted to off-the-shelf medicines and devices.

### What is this preparation for?

---

This is a custom-formulated preparation designed to better meet your individual needs. This practice's desire is to help you achieve better outcomes. Its active ingredients will work together toward this goal.

**Azelaic Acid** is a saturated acid that has demonstrated efficacy in stopping the growth of skin bacteria that causes acne. **Azelaic Acid's** antimicrobial action may be attributable to inhibition of microbial cellular protein synthesis.

**Niacinamide** also known as **Nicotinic Acid** is a B vitamin, specifically **Vitamin B3**. **Niacinamide** is touted in Rx dermatology and skincare products for reducing redness and irritation. **Niacinamide** has antioxidant and anti-inflammatory properties that complement the cell communicating role. These properties make **Niacinamide** an excellent conditioning agent that reduces the harshness of the additional active ingredients contained within the formula.

### How to use this preparation?

---

Use this preparation exactly as your doctor instructed you.

Additional directions:

---

---

### How to store this preparation

---

This preparation should be stored in a cool, dry and dark place.

**Warning:** You may have an allergic reaction to any of the ingredients in this custom-formulated preparation. It is advisable that you first test the preparation before applying it on a large area of the body. Apply a very small amount of cream/lotion/gel/solution on a quarter size area on your forearm to test. Let it stay for about 30 minutes. If you have a fast reaction type of allergy, the area will be red and itchy. If you have a slow reaction type of allergy, the same symptoms may occur after 48 hours. If any of these situations occur, stop using the preparation and call your health care practitioner. The absence of any reaction to this test is no guarantee that an allergic reaction may not occur during future uses of this preparation.