

# Colonoscopy Bowel Prep Instructions

---

## SUPREP

### Today

- Take time off of work for the day of your colonoscopy.
  - Find a responsible person to either drive you home or ride with you in a car service like Uber after your colonoscopy. **You will not be able to drive yourself after the procedure.**
  - If you are taking any blood thinner medications other than aspirin or NSAIDs**, please contact the doctor who prescribes this medication. Depending on the specific blood thinner, these will typically need to be stopped between 2-5 days before your procedure.
  - If you take medications for diabetes or weight loss**, please contact the doctor who prescribes this medication. Depending on the specific medication, this may need to be stopped altogether between 1-7 days before your procedure, or the dose may only need to be reduced starting the day before your colonoscopy.
- 

### 7 Days Before Your Colonoscopy

- Go to your pharmacy and buy:**
    - SUPREP** (this is a prescription medication)
    - Baby Wipes**, this can be more soothing than toilet paper
    - If you typically experience constipation, then buy 1 bottle of **Milk of Magnesia**
  - Stop** taking iron pills, multivitamins with iron, and fiber supplements (Metamucil, Citrucel, and Benefiber).
  - Make sure a plan is finalized if you need to stop taking blood thinners, diabetes medications, or weight loss medications prior to your colonoscopy.
- 

### 5 Days Before Your Colonoscopy

- Follow a low fiber diet:** avoid whole grain breads, nuts, seeds, popcorn, corn, fruits, and vegetables for **5 days** before you begin your colonoscopy prep.
- 

### 2 Days Before Your Colonoscopy

- Remain well hydrated. Drink at least 8 glasses of water today.
  - If you typically have constipation, then take 2 tablespoons of milk of magnesia at 7 PM.
- 

### The Day Before Your Colonoscopy

- At 8 AM** eat a light breakfast (i.e. toast, eggs, cereal without nuts or whole grains)
  - For the remainder of the day, follow a strict clear liquid diet. Do not eat any additional food.**
    - Clear liquids are: water, apple juice, clear broths, Jell-O, black coffee or tea without milk/cream, Sprite, Ginger ale, Gatorade, popsicles (no sherbet or fruit bars), flavored ice. **Nothing red or purple.**
  - Make sure you are drinking clear liquids throughout the day to remain well hydrated.
  - At 5 PM**, pour one 6 oz bottle of SUPREP into a container and fill with cold water to the 16 oz line. Mix these together and drink. Then drink an **additional** two 16 oz containers of water over the next hour. If feeling sick to your stomach, take breaks or drink other clear liquids to settle your stomach.
- 

### The Day of Your Appointment

- 6 hours before your arrival time**, pour the remaining 6 oz bottle of SUPREP into a container and fill with cold water to the 16 oz line. Mix these together and drink. Then drink an **additional** two 16 oz containers of water over the next hour. (Depending on the time of your procedure, you may need to start drinking this in the middle of the night.) You must finish this at least **3 hours before** your colonoscopy. **This half is the most important part of the bowel prep!**
- Take your usual medications with a few sips of water unless you are told otherwise.
- Starting **3 hours** before your colonoscopy **do NOT have anything else to drink.**
- Be sure to check the location of your procedure listed above. We look forward to seeing you!