

AGE	IMMUNIZATION	DISEASE PROTECTION
Newborn	Hepatitis B	Hepatitis B
1 Month	Hepatitis B	Hepatitis B
2 Months	Pprevnar 13, Pentacel, Rotavirus (oral)	Pneumococcal disease Polio Diphtheria Tetanus Whooping cough (pertussis) Hemophilus Influenza type B Rotavirus
4 Months	Pprevnar 13, Pentacel, Rotavirus (oral)	Pneumococcal disease Polio Diphtheria Tetanus Whooping cough (pertussis) Hemophilus Influenza type B Rotavirus
6 Months	Pprevnar 13, Pentacel, Rotavirus (oral)	Pneumococcal disease Polio Diphtheria Tetanus Whooping cough (pertussis) Hemophilus Influenza type B Rotavirus
9 Months	Hep B	Hepatitis B
12 Months	MMR, Varivax, Hep A	Measles / Mumps / Rubella Varicella (Chicken pox) Hepatitis A
15 Months	Pprevnar 13, Pentacel	Pneumococcal disease Polio Diphtheria Tetanus Whooping cough (pertussis) Hib Influenza
18 Months	Hep A	Hepatitis A
2 Years.	Vaccines given if behind	
2½ Years	Vaccines given if behind	
3 Years.	Vaccines given if behind	

4 Years	MMR, Varivax (DTaP, Polio)	Measles / Mumps / Rubella Varicella (Chicken pox)
5 Years	DTaP, IPV	Diphtheria Tetanus Whooping cough (pertussis) Polio
Annual wellness visits	Vaccines given if behind	
10 Years	TDaP	Diphtheria Tetanus Whooping cough (pertussis)
11 Years	Menactra, HPV	Meningitis (4 strains) Human papilloma virus
11 - 12 years	HPV	Human papilloma virus
Annual wellness Visits	Vaccines given if behind	
16 Years	Menactra	Meningitis (4 strains)
16 - 18 years	May have Bexsero May have Td booster	Meningitis B Tetanus and diphtheria

In addition to the vaccines listed here, an annual influenza vaccine is also recommended by our office as the best protection against seasonal flu. This vaccine is best administered prior to the influenza season - typically September through November.