










**COLONOSCOPY: TAKING YOUR DIABETES MEDICATION(S)  
DAY BEFORE PROCEDURE**

*Follow the clear liquid diet and adjust your medications*

<b>If you take:</b>	<b>Breakfast</b>	<b>Lunch/Dinner</b>	<b>Bedtime</b>
Actoplus, Actos, Avandamet, Avandia, Januvia, Janumet, Met, Metformin	Take your usual dose	Take your usual dose	
Amaryl, Avandaryl, Duetact, Glipizide, Glucovance, Glyburide, Metglip, Prandin, Starlix	DO NOT TAKE	DO NOT TAKE	
Lantus or NPH Insulin	Take your usual dose		Take your usual dose
Humalog, Novolog or Regular Insulin	Take ½ the usual amount if you take a fixed dose <b>OR</b> Cover your carbs with your usual carb ratio	Take ½ the usual amount if you take a fixed dose <b>OR</b> Cover your carbs with your usual carb ratio	
Novalog Mix 70/30, Novolin 70/30, Premixed Insulin 75/25	Take ½ the usual dose at breakfast	Take ½ the usual dose at dinner	

**DAY OF PROCEDURE**

*Do not make up or “double up” on any missed medications after the procedure*

<b>If you take:</b>	<b>Breakfast</b>	<b>Lunch/Dinner</b>	<b>Bedtime</b>
Actoplus, Actos, Avandamet, Avandia, Januvia, Janumet, Met, Metformin	DO NOT TAKE	DO NOT TAKE	
Amaryl, Avandaryl, Duetact, Glipizide, Glucovance, Glyburide, Metglip, Prandin, Starlix	DO NOT TAKE	Resume your usual dose if eating	
Lantus or NPH Insulin	Take ½ of the usual dose		Resume you usual dose
Humalog, Novolog or Regular Insulin	DO NOT TAKE	Resume your usual dose if eating	
Novalog Mix 70/30, Novolin 70/30, Premixed Insulin 75/25	DO NOT TAKE	Resume your usual dose if eating	