



# Bowel Preparation Instructions - PLENVU Any questions please call (631)-289-0300 ext. 224.

#### 3 DAYS PRIOR to the PROCEDURE:

- Do not eat foods high in fiber.
- Examples are whole grain breads and cereals, fruits, nuts, seeds, quinoa, popcorn, and cooked or raw vegetables.
- Do not take any fiber supplements.

#### On the DAY BEFORE Your PROCEDURE:

#### **DIET**

- Consume only clear liquids the entire day before
- Okay to consume one protein shake, before 11 am, but only from the following lists:
  - Protein powder mixed with water or milk
  - Pre-made Protein shake
  - AVOID FRUIT or VEGETABLE shakes.

At 11 AM: Begin Drinking ONLY clear liquids. NO MORE PROTEIN SHAKES.

A clear liquid is one that can you see through. Examples of clear liquid are as follows:

- Clear beef or chicken broth
- Popsicles, except for red, purple or blue
- Coconut water (NOT coconut milk)
- Strained fruit juice such as apple or white grape juice
- Coffee and tea (without cream or dairy products)
- Gatorade
- Soft drinks
- Kool-Aid
- Jell-O
- Do not drink alcohol

# PREP - Procedure before 9am

#### 7pm - Start 1st Dose:

- 1. Use the mixing container to mix the contents of Dose 1 pouch with at least 16 ounces of water by shaking or using a spoon until it's completely dissolved. This may take up to 2-3 minutes. Take your time slowly finish the dose within 30 minutes.
- 2. Refill the container with at least 16 ounces of clear liquid. Again, take your time and slowly finish all of it within 60 minutes.

#### 9pm - start 2nd dose

- Follow steps 1-2 above.
- Starting at midnight, you may have nothing by mouth, including water
- Drink plenty of clear liquids during and after bowel-prep process to help prevent dehydration

#### On the Morning/Day of your PROCEDURE

- If you take daily medication(s), you may do so up until 2 hours before your scheduled report time with small sips of water.
- If you take blood thinners, follow the instructions your health care provider gives you.
- Arrange for a ride home. You will be drowsy after the exam and will need an escort, 18 years or older. If taking a medical cab service, you will need an escort as well to take you home.

# The importance of a GOOD PREP

Your prep is one of the most important parts of your colonoscopy. Without a good prep, your bowel empty and clean, your gastroenterologist will not be able to see your colon clearly. This could lead to missing a polyp/lesions or rescheduling your colonoscopy.

#### PREP - Procedure 9am or later

# 7pm - Start 1st Dose:

- 1. Use the mixing container to mix the contents of Dose 1 pouch with at least 16 ounces of water by shaking or using a spoon until it's completely dissolved. This may take up to 2-3 minutes. Take your time slowly finish the dose within 30 minutes.
- 2. Refill the container with at least 16 ounces of clear liquid. Again, take your time and slowly finish all of it within 60 minutes.
- Drink plenty of clear liquids during and after bowel-prep process to help prevent dehydration

# On the Morning/Day of your PROCEDURE

#### 6am – start 2nd Dose:

- Follow steps 1-2 above.
- Be sure to finish drinking all of your laxative by 7am.
- If you take daily medication(s), you may do so up until 2 hours before your scheduled report time with small sips of water.
- If you take blood thinners, follow the instructions your health care provider gives you.
- Arrange for a ride home. You will be drowsy after the exam and will need an escort, 18 years or older. If taking a medical cab service, you will need an escort as well to take you home.

# The importance of a GOOD PREP

Your prep is one of the most important parts of your colonoscopy. Without a good prep, your bowel empty and clean, your gastroenterologist will not be able to see your colon clearly. This could lead to missing a polyp/lesions or rescheduling your colonoscopy.