

**Data regarding the varying improvement  
of Voiding Symptoms with the  
continuous use of the Peenuts®  
Formula of 6-12 months or longer**

MILD SYMPTOMS (1-7)			MODERATE SYMPTOMS (8-19)			SEVERE SYMPTOMS (20-35)		
Pre-Peenuts	Post-Peenuts	% Change	Pre-Peenuts	Post-Peenuts	% Change	Pre-Peenuts	Post-Peenuts	% Change
1.5	0.5	67%	16.5	4	76%	20.5	9	56%
4	0.5	88%	13.5	4.5	67%	20	11.5	43%
2.5	0.5	80%	15.5	5.5	65%	30.5	13.5	56%
7.5	3.5	53%	8.5	0.5	94%	20.5	10	51%
7.5	3.5	53%	13	2	85%	20.5	8.5	59%
6	1	83%	12.5	2	84%	21	0.5	98%
5.5	0	100%	10	2.5	75%	21.5	8	63%
4.5	2	56%	10.5	1.5	86%	23.5	16.5	30%
5.5	2.5	55%	11	1.5	86%	25	16	36%
6.5	2	69%	13	2	85%	23	10.5	55%
4	0	100%	15	1.5	90%	22.5	10	56%
5.5	2.5	55%	11.5	1.5	87%	25.5	15.5	40%
7.5	1.5	80%	18.5	5.5	70%	20	4.5	78%
7	0	100%	5.5	0	100%	22	15.5	30%
5	2	60%	16.5	8.5	48%	21.5	2	91%
3.5	0.5	86%	11	2.5	77%	21	4.5	79%
2.5	2	20%	17	1.5	91%	23.5	12	49%
6	3	50%	11.5	4.5	61%	28.5	12.5	57%
7.5	2	73%	14	2.5	82%	20.5	7	66%
7	0.5	93%	11.5	6.5	43%	23.5	10	57%
5.5	0	100%	9.5	1.5	84%	20	7	65%

MILD SYMPTOMS (1-7)			MODERATE SYMPTOMS (8-19)			SEVERE SYMPTOMS (20-35)		
Pre-Peenuts	Post-Peenuts	% Change	Pre-Peenuts	Post-Peenuts	% Change	Pre-Peenuts	Post-Peenuts	% Change
			8	1.5	81%	24.5	16	34%
<b>N=21</b>		<b>72%</b>	14	6.5	53%	25	4.5	86%
		<b>Improved</b>	14	5	64%	22	5	77%
			12	3.5	71%	25.5	3.5	86%
			15	5	67%			
			19.5	4	80%	<b>N=25</b>		<b>60%</b>
			9.5	3.5	63%			<b>Improved</b>
			18.5	10	46%			
			16	5	69%			
			9.5	2.5	74%			
			16.5	8	52%			
			20	7	65%			
			10.5	3.5	67%			
			11.5	1.5	87%			
			14.5	4.5	69%			
			19.5	8.5	56%			
			10.5	6	43%			
			11.5	7.5	35%			
			11	7	36%			
			9.5	2	79%			
			15.5	10	35%			

MILD SYMPTOMS (1-7)			MODERATE SYMPTOMS (8-19)			SEVERE SYMPTOMS (20-35)		
Pre-Peenuts	Post-Peenuts	% Change	Pre-Peenuts	Post-Peenuts	% Change	Pre-Peenuts	Post-Peenuts	% Change
			8.5	1.5	82%			
			15.5	5.5	65%			
			14.5	10.5	28%			
			12.5	1	92%			
			11.5	3	74%			
			13.5	7.5	44%			
			15.5	8.5	45%			
			14	2.5	82%			
			16.5	11.5	70%			
			10	3.5	65%			
			15.5	5	68%			
			15	3	80%			
			15.5	5.5	65%			
			13.5	4	71%			
			15.5	7	55%			
			15	1	94%			
			14.5	4.5	69%			
			16	2.5	85%			
			9	5	45%			
			17	4.5	74%			
			9.5	2.5	74%			

MILD SYMPTOMS (1-7)			MODERATE SYMPTOMS (8-19)			SEVERE SYMPTOMS (20-35)		
Pre-Peenuts	Post-Peenuts	% Change	Pre-Peenuts	Post-Peenuts	% Change	Pre-Peenuts	Post-Peenuts	% Change
			17	3.5	79%			
			12	7.5	38%			
			16	5.5	67%			
			13.5	3.5	74%			
			16.5	7.5	55%			
			14.5	0.5	97%			
			11	2.5	78%			
			8.5	0.5	95%			
			11.5	3.5	70%			
			19	6	68%			
			10.5	2.5	76%			
			18.5	11.5	38%			
			17.5	5.5	69%			
			16.5	8.5	48%			
			16.5	6.5	61%			
			15.5	5	61%			
			9	2	78%			
			10.5	0	100%			
			17	1.5	91%			
			11.5	4.5	61%			
			12.5	2	84%			

MILD SYMPTOMS (1-7)			MODERATE SYMPTOMS (8-19)			SEVERE SYMPTOMS (20-35)		
Pre-Peenuts	Post-Peenuts	% Change	Pre-Peenuts	Post-Peenuts	% Change	Pre-Peenuts	Post-Peenuts	% Change
			14	2.5	82%			
			18.5	4	79%			
			17	10.5	39%			
			15.5	7	55%			
			17	7	59%			
			17	9	47%			
			16.5	8.5	49%			
			14	2	86%			
			13.5	4	71%			
			8.5	3.5	59%			
			11.5	3.5	70%			
			15	2	87%			
			17.5	5	72%			
			10	3	70%			
			14	3	79%			
			10	5	50%			
			17	5.5	68%			
			13	7.5	43%			
			9	1	89%			
			17.5	7	60%			
			17	3	83%			

MILD SYMPTOMS (1-7)		MODERATE SYMPTOMS (8-19)		SEVERE SYMPTOMS (20-35)	
Pre-Peenuts	Post-Peenuts	Pre-Peenuts	Post-Peenuts	Pre-Peenuts	Post-Peenuts
	% Change		% Change		% Change
		17	6.5		
		16	6		
		18.5	7.5		
		12	6		
		11.5	3.5		
		13.5	2.5		
		12	2.5		
		11	5		
		16	3.5		
		11	3		
		18.5	12.5		
		15	3		
		15.5	5.5		
		13.5	4		
		15.5	7		
		15	1		
		14.5	4.5		
		16	2.5		
		9	5		
		17	4.5		
		9.5	2.5		

MILD SYMPTOMS (1-7)		MODERATE SYMPTOMS (8-19)		SEVERE SYMPTOMS (20-35)	
Pre-Peanuts	Post-Peanuts	Pre-Peanuts	Post-Peanuts	Pre-Peanuts	Post-Peanuts
% Change		% Change		% Change	
	17	3.5	74%		
	12	7.5	38%		
	16.5	5.5	67%		
	13.5	3.5	74%		
	16.5	7.5	55%		
	14.5	0.5	97%		
	11	2.5	78%		
	8.5	0.5	95%		
	11.5	3.5	70%		
	19	6	68%		
	10.5	2.5	76%		
	18.5	11.5	38%		
	17.5	5.5	69%		
	16.5	8.5	48%		
	16.5	6.5	61%		
	9	2	78%		
	10.5	0	100%		
	17	1.5	91%		
	11.5	4.5	61%		
	12.5	2	84%		
	14	2.5	82%		



MILD SYMPTOMS (1-7)			MODERATE SYMPTOMS (8-19)			SEVERE SYMPTOMS (20-35)		
Pre-Peenuts	Post-Peenuts	% Change	Pre-Peenuts	Post-Peenuts	% Change	Pre-Peenuts	Post-Peenuts	% Change
			16.5	4	76%			
			13.5	4.5	67%			
			15.5	5.5	65%			
			8.5	0.5	94%			
			13	2	85%			
			12.5	2	84%			
			10	2.5	75%			
			10.5	1.5	86%			
			11	1.5	86%			
			13	2	85%			
			15	1.5	90%			
			11.5	1.5	87%			
			18.5	5.5	70%			
			16.5	8.5	48%			
			11	2.5	77%			

N=162

70% Improvement