


SUTAB

follow the instructions on this paper DO NOT FOLLOW THE INSTRUCTIONS ON THE SUTAB BOX

PLEASE READ ALL INSTRUCTIONS CAREFULLY WHEN PROCEDURE IS SCHEDULED

7 DAYS PRIOR	3 DAYS PRIOR	ONE DAY PRIOR	THE DAY OF PROCEDURE
<p>Supplies</p> <ul style="list-style-type: none"> • SUTAB Bowel Prep Kit and Ondansetron Prescription– These will be sent to Gifthealth Pharmacy (please call 833-614-4438 to coordinate receiving these if you haven't already). • Purchase Gas-X tablets (Simethicone). The cherry flavor, pink tablets are approved to use during the prep.  <p>If you take medication to thin your blood, such as Coumadin (warfarin), Plavix (clopidogrel), Xarelto (rivaroxaban) or Pradaxa (dabigatran), Eliquis (apixaban), Aggrenox (aspirin/dipyridamole), etc., ask the doctor that prescribed it for instructions prior to stopping the medication. Aspirin may be continued.</p> <p>STOP all fiber supplements or medications containing iron 7 days before your appointment. This includes multi-vitamins with iron, Metamucil, Citrucel and Fibercon.</p> <p>Arrange for a driver on the day of your procedure.</p>	<p>STOP consuming all high fiber foods/roughage.</p> <ul style="list-style-type: none"> • NO raw vegetables • NO corn (raw or cooked) • NO whole wheat or high fiber breads • NO nuts or popcorn • NO bran or bulking agents <p>Confirm that you have a driver for the day of the procedure.</p> <p>A taxi, Uber, or bus are NOT allowed unless you are accompanied by a family member or friend.</p> <p>Review paperwork and sign</p> <ul style="list-style-type: none"> • Multi-authorization • Notice of Privacy Practices 	<p>NO SOLID FOOD NO ALCOHOL</p> <p>Clear liquids ALL DAY</p> <p>AVOID anything red or purple in color.</p> <p>NO milk products or non-dairy creamer.</p> <p>SEE ATTACHED LIST OF CLEAR LIQUIDS</p> <p>DIABETICS: Do NOT take oral medication DIABETA, GLUCOPHAGE, METFORMIN or JANUVIA</p> <p>Take ½ of your regular insulin dose</p> <hr/> <ul style="list-style-type: none"> • STEP 1: At 3:00 pm take 2 Ondansetron tablets • STEP 2: At 4:00pm Fill the plastic container that comes with the kit with water (there is a line on the container). Drink the entire cup over the next hour. • STEP 3: AT 5 PM Fill the plastic container that comes with the kit with water again (there is a line on the container). Open one of the two bottles in the kit. It will contain 12 tablets. Swallow one tablet at a time with a sip of water every 3-4 minutes. If you run out of water, you can drink more water. If you become uncomfortable or nauseated take a break for a few minutes then resume taking tablets every 3-4 minutes until the 12 are gone. Take one Gas-X tablet. This can take about an hour. • STEP 4: About one hour after taking the last tablet, fill the container with water again and drink over 30 minutes. • STEP 5: About 30 minutes after finishing the second container of water, fill the container again with water and drink that over 30 minutes. <p>• Note: We are all different, some people will have a rapid response with bowel movements shortly after beginning the preparation while others will take longer. If you experience preparation-related symptoms (for example, nausea, bloating or cramping), pause or slow the rate of drinking the additional water until your symptoms diminish.</p> <ul style="list-style-type: none"> • You may continue clear liquids 	<p>NO SOLID FOOD NO ALCOHOL</p> <p>DIABETICS: Do NOT take oral medication DIABETA, GLUCOPHAGE. METFORMIN or JANUVIA</p> <ul style="list-style-type: none"> - If you are Insulin dependent, do not take your morning dose of insulin. - You must check your blood sugar before leaving home. If your level is above 300, please take your Regular insulin according to your sliding scale. - Please bring your insulin with you to procedure center. <p>All patients may take morning medications with sips of water.</p> <ul style="list-style-type: none"> • At _____ (7 hours before your procedure time) fill the clear container up to the line with water and drink over the next hour • At _____ (6 hours before your procedure)Take 2 Gas-X tablets. Repeat STEPS 3 through 5. Be sure to wait 3-4 minutes between pills. • It is important to have completed all these steps at least 4 hours before your colonoscopy. <p>You may continue to drink water until: 2 HOURS PRIOR to procedure.</p> <p>Drinking liquids within these 3 hours will cause us to cancel or postpone your procedure. **Also no gum, hard candy, mints and tobacco products in these 3 hours.</p> <p>If you are unsure of your colon cleansing, please call the office for assistance: Akron Digestive Disease Consultants 330-869-0124 The Gastroenterology Group 330-869-0954</p>

A Colonoscopy has been recommended to you. This examination involves guiding a tubular, flexible instrument through the rectum and into the entire colon. The preparation for the procedure is IMPORTANT. It does cause diarrhea and possibly abdominal cramping.

CLEAR LIQUIDS – No RED or PURPLE

Must be something you can see through...



- Gatorade or PowerAde
- Clear broth or bouillon – chicken or beef
- Coffee or Tea (*no milk or non-dairy creamer*)
- Carbonated and Non-Carbonated Soft Drinks
- Kool-Aid or Crystal Light
- Strained Fruit Juices (*no pulp*)
- Jell-O, Popsicles, or Italian Ice

DON'T FORGET - these foods are **NOT ALLOWED!**



- Milk or milkshakes
- Cream or non-dairy creamer
- Orange, Grapefruit or Tomato Juice
- Creamed Soups or any soup other than broth
- Oatmeal
- Cream of Wheat

Colon Cleansing Tips

- ➡ You may use over the counter hydrocortisone creams, tucks pads or baby wipes as necessary for skin irritation. Be sure to get alcohol free!
- ➡ Chill the solution in refrigerator or by placing in a bowl of ice. DO NOT add ice to the solution.
- ➡ Stay near the toilet! You will have diarrhea, which can be quite sudden...this is normal!
- ➡ If you experience nausea or vomiting, rinse your mouth with water and take a break 😊 Wait 45 minutes and then resume drinking the prep. If necessary, slow down, let your stomach settle between glasses.
- ➡ It is common to experience abdominal discomfort until the stool has flushed from your colon.
- ➡ Some find it easier to drink through a straw.
- ➡ Eventually your stools will be liquid, and clear enough to see the bottom of the toilet.
- ➡ If you feel for any reason your prep is not working, please be sure to call us so we can assist you in completing the prep process.
- ➡ **EVEN AFTER STOOLS BECOME CLEAR YELLOW LIQUID – YOU MUST COMPLETE ALL STEPS OF THE PREP!**

THINGS TO BRING WITH YOU!

- ✓ A RESPONSIBLE DRIVER
- ✓ YOUR INSURANCE CARDS
- ✓ A PHOTO ID
- ✓ SIGNED PAPERWORK FOR FACILITY